

Rifting systems and its significance for hydrocarbon exploration in the Netherlands

F.F.N. van Hulten

Energie Beheer Nederland B.V. - P.O. Box 6500, 6401JH - Heerlen, The Netherlands

Abstract

Rifting has shaped the Dutch subsurface in significant way. A brief summary of general knowledge related to this topic is given as a background to the symposium on rifting systems and its significance for hydrocarbon exploration in the Netherlands. This symposium coincides with increased interest in hydrocarbon exploration in the Netherlands. New data becoming available is creating new insights in mature exploration areas. The improved subsurface imaging of geological structures, with better 3D acquisition and computer aided imaging tools, makes it increasingly clear that trapping and accumulation of hydrocarbons in the Netherlands can be much more complex than earlier realized. The North Sea rifting phenomena are in a complicated way connected to the Upper Rhine Graben via the Netherlands. Also the non geologist in the country is periodically reminded of the rifting process because of minor earthquakes.

Basins related to rifting play an important role in the spectrum of basin types. Rifting is widely recognized as an important factor that has shaped our oil and gas fields and has been the subject of numerous papers describing rift phenomena in the country and the wider Southern North Sea area. The importance of structuration caused by the rifting process for oil and gas exploration is well documented for selected cases, but the subject is somewhat neglected compared to the numerous papers available in the Northern North Sea area of the UK and Norway.

In the following a short anthology of topics related to rifting and related structuration is given and its relevance for exploration in the Netherlands is briefly reviewed.

1.0 Introduction

The North Sea is generally seen as an example of a rifted basin. The Netherlands is an integral part of this basin and forms the complex transition of the North Sea rift system to the Cenozoic Rhine Graben system, part of the European Cenozoic rift system (Sissingh, 2006a). Rifting was already described decennia before continental drift was accepted but the new global tectonics theory that including major plate movements after 1967 gave rifting a different meaning. Present day examples of presently active rifts are the East African Rift and the mainly German Cenozoic Rhine Graben rift. The Dutch Roer Valley Graben, the active present-day rift basin of the Netherlands, is the northerly extension of the latter. The first description of these areas goes back to the first part of the twentieth century but the full understanding of all rifting processes and all its related structuration is still developing and subject of this workshop.

Dr. P.A. Ziegler has been very important to promote the rift concept in the Netherlands but in a mail to the conveners of this conference he writes: “...*rifting was invented much before my active time. Just think of Hans Cloos (1936) as an early rifter! My personal work was much later and centered on understanding the evolution of the Arctic-North Atlantic and Tethys rift systems and also the evolution of the European Cenozoic rift system. That the Permian Oslo, the Mesozoic North Sea and the Cenozoic Rhine-Rhône rift systems have nothing to do with each other should by now be clear to anybody who had freed himself from Stille's misbegotten Mittelmeer-Miöse rift concept and read my publications.*”

2.0 Rifting definition

Rifting is a fundamental geological process and many descriptions can be found. A widely used handbook like Kearey & Vine (1990) rifts has proposed the following:

Rifts are defined as elongate depressions where the entire thickness of the lithosphere has deformed under the influence of tensional forces.

When this description is compared to current internet definitions one can see an increasing emphasis on continental plate processes.

Wikipedia:

In geology, a **rift** is a place where the Earth's crust and lithosphere are being pulled apart. Typical features are a central linear down dropped fault segment, called a graben, with parallel normal faulting and rift-flank uplifts on either side forming a rift valley. The axis of the rift area commonly contains volcanic rocks and active volcanism is a part of many but not all active rift systems. Rifts are distinct from Mid-ocean ridges, where new oceanic crust and lithosphere is created by seafloor spreading. In rifts, no crust or lithosphere is produced. If rifting continues, eventually a mid-ocean ridge may form, marking a divergent boundary between two tectonic plates.

Indopedia:

In geology, a **rift** is a place where the Earth's lithosphere is expanding. Typical features are a central linear down dropped fault segment, called a graben, with parallel normal faulting on either side. The central portion of the rift area commonly contains volcanic rocks and active volcanism is a part of active rift systems. Rifts are located at **divergent boundaries** between two tectonic plates. Between oceanic plates the rifts occur as oceanic ridges and where these oceanic ridges intersect continental crust rift valleys result as the continent begins to split.

Rifting is presently not restricted to the depression only in agreement with Ziegler (2001) describing the process as a combination of far-field plate boundary stresses and frictional forces exerted on the base of the lithosphere by the convecting upper mantle. This implies that the rifting process looks at a wider (plate) area and beyond the extensional phase. Such an approach has been used for the area south of the Netherlands (Sissingh, 2006a, 2006b).

2.1 Some general characteristics of rifting

Keary & Vine (1990) give a listing of a number of characteristics of rifting that give clear insight:

- The term rift applies to major lithospheric features and does not encompass small scale structures.
- Geological studies demonstrate that the bounding faults of rift valleys are of normal type. Rifts consequently develop in response to tensional stresses but there are theories that note that compression plays a role.
- The location of rifts is often controlled by crustal weakness
- The lithosphere underlying rifts is generally abnormally thin and invaded by high low velocity, low density and high temperature material
- Horizontal displacement is relatively small [8-10 km for the Rhine Graben]
- Rifts are spatially associated with **domal uplifts**.
- Rifts may form an interlinked network dividing a continental plate
- Rifts contain often contain extensive **volcanic rocks**. Where present these rocks are mildly to strongly alkaline and the alkalinity decrease with time and increases with distance to the rift axis.

2.2 Classification of rifting

It is customary to classify rifts. *One of the questions that we like to pose at this workshop is if we are in an active or paleo rift area.*

- Rifting associated with “Break Up” is active rifting
 - Precursor of a new ocean
 - Splitting creates a number of not utilized rifts (Aulacogens)
- Continental collision creates compression and areas distant from the suture can develop a tensional regime (Rhine Graben)
- Distinction is made between rifts generated by continental rupture or collision (mantle activated).
- Lithosphere activated (passive) rifts are not activated before rupture by upwelling asthenosphere.
- Mantle activated rifts are characterized by large volumes of extrusive rocks.

2.3 Structure of continental rifts

Continental rifts are bounded by normal faults. Rifts develop in response to tensional stresses.

Mega structures related to rifts are:

- Transcurrent faults
- Strike slip faults
- Pull apart basins
- Triple junctions
- Aulacogens [Central Graben is an example (Kearey & Vine, 1990)]

2.3.1 Crust and rifting

Seismic refraction shows that crust underneath rifts is significantly thinner than beneath adjacent areas. This has also been noted in the Netherlands. Deep seismic shows a thinner crust in the Roer Valley Graben area (Rijkers & Duin, 1994).

Typical thickness of the continental crust is on average 35 km (Kearey & Vine, 1990) and less than 20 km in active (rifting) areas and up to 80 km beneath young fold belts. Observations in the Netherlands confirm this with a thickness of about 38 km underneath the massifs (London Brabant Massif). In the rifted areas the thickness is thinner. Over a limited area underneath the Roer Valley Graben the depth of the Moho is 28 km. The location of the earthquake of 1992 observed at a depth of about 18 km is in line with the observation of the deep seismic (Duin et al., 1994) and is related to one of the boundary faults.

It is interesting to note that a large part of the northern offshore also has a thinner crust (Duin et al., 1994; Abramovitz & Thybo, 2000). Danish deep seismic lines (MONA LISA -1,-2) suggest that the Avalonian basement rock can be divided in a northern area with thinner crust and southern area (in the Netherlands a line parallel to the Frisian islands), that displays a thicker with a good developed middle crust (Abramovitz et al., 1999). The northern thinner crust is possibly oceanic crust (including meta-sedimentary cover), with a basement velocity that is relative low (5.8-5.9 km/s) compared to the southern area with blocks that share cratonic properties. The lithosphere in the northern offshore is therefore possibly weaker and more prone to rifting vs. the southern part of the Netherlands that consist of an amalgamation of different terranes during the Caledonian orogeny (Abramovitz et al., 1999) with suspected Caledonian blocks with a higher lithospheric strength like the Texel IJsselmeer High and the Zandvoort-Maasbommel-Kreveld High.

Gravity surveys (Visser et al., 1987) have shown that in a late tectonic sense the uplifted regions are in isostatic equilibrium; however negative Bouguer anomalies are locally present

over the rifted Roer Valley Graben structure. This graben, that is active presently, is characterized by a narrow 30 km wide and over 100 km wide anomaly. Onshore an anomaly is also noticeable over the Lauwerszee Trough.

3.0 Rifting in the Netherlands

For a good introduction for the structural history of the Netherlands can be referred to Heybroek (1974) and van Wijhe (1987). Recently an update was published by De Jager (2007). Regional structure maps of the Netherlands can be found in Duin et al. (2006).

3.1 Hydrocarbon discoveries in rift areas of the Netherlands

Geological description of graben tectonics in the Netherlands started in the Roer Valley Graben area and is related to coal exploration and exploitation in the first part of the twentieth century in the southern part of the country. For example reverse movements were already noted by W.C. Klein in 1910 in South Limburg (Heybroek, 1967).

The location of this rift valley as other tectonic elements can be found and are described in Adrichem Boogaert & Kouwe (1997). The predecessor of this graben to the west, in what is called the West Netherlands Basin, yielded several oil and gas discoveries starting in 1938 (the first oil) or 1944 (the first field). See for the oil and gas fields of the West Netherlands Basin Bodenhausen & Ott (1981), De Jager et al. 1996 and Alvaro Racero-Baena & Drake (1996). Exploration in the onshore part of the basin was important in the 1950's and 1960's, but continues till today. The first oil offshore was found by Tenneco in 1970 in F18 in the Dutch Central Graben. A flow of 325 m³ of light (31 API°) oil was measured during flow tests from Jurassic sandstones at a depth of 2,500 m but was not commercial. Later a number of significant oil and gas finds were made in the Dutch Central Graben. The F3-FB Jurassic condensate/oil discovery was a major find in 1994. The southern rift basins yielded several oilfields: Kotter (De Jong & Laker, 1992), Logger (Goh, 1996), Helm (Hastings et al., 1991), Helder/Hoorn (Roelofsen & de Boer, 1991) and the Rijn (Alberts et al., 2003) oil field in P15.

Rifting processes affect not the basin areas only but are of importance for a much larger area including the Lower Saxony Basin, the Lauwerszee Trough. Of particular interest are the Broad Fourteens Basin and the adjacent area because many of the Dutch offshore Rotliegend gas fields can be found at its northern edge.

3.2 Geological history and rifting in the Netherlands

Rifts play an important role in the development of sedimentary basins of the Netherlands. Several rift cycles have been described in the country in a number of different megatectonic settings. The duration of the rifting is highly variable.

3.2.1 The earliest rifting

The geological history of the Netherlands more or less starts with rifting. De Jager, (2007) state that in the late Ordovician the Avalonia micro plate rifted away from Gondwana and joined Baltica in the late Ordovician with the closure of the Tornquist Ocean. The resulting collage with blocks like the London Brabant Massif and the Texel IJsselmeer High influence the development of rifting. The rupturing of the Jurassic rifting seems to halt against these Caledonian blocks.

The next period where rifting may be an important structural factor is during the Devonian into the Mississippian. Fraser and Gawthorpe suggest that (back arc) rifting initiated a series of half grabens. The Campine Basin in the Southern Netherlands is one of these half grabens. Also Ziegler (1982) noted already that before the late Carboniferous (Variscan) compressional events start, Dinantian half grabens can be explained by Carboniferous rifting (Arctic – north Atlantic

rift system). Kockel (2002) suggests a similar explanation and adds that part of this rifting has been obscured by the later Variscan events.

After the Variscan events there is little evidence for rifting in the Netherlands contrary to reports from Germany (Ziegler, 1990). Kockel (2002) also notes that for block faulting older than Upper Rotliegend no clear major faults or grabens can be outlined. In the Upper Rotliegend the Schneverdingen sands are deposited in block faulted grabens. In the Netherlands thickness differences of deposits of Rotliegend and Zechstein age do not indicate rifting grabens. The nature of the depocenters of the Permian basin (Geluk, 2005), that is so clearly expressed in their facies patterns, is explained as a result of the decay of the thermal anomaly introduced during the Permo-Carboniferous (Van Wees et al., 2000).

3.2.2 The breaking up of Pangea

In the Mesozoic a number of rifting events can be distinguished that are related to the break up of Pangea. The opening of the Atlantic continues in the Tertiary. Because of the Alpine collision partly due to the opening of the Southern Atlantic a number of compressional events change the stress field over large areas and periods with significant inversion are more dominant than periods with extension (Ziegler, 1982; Ziegler, 1990).

3.2.2.1 Triassic rifting

The late Triassic rifting is also known as Early Kimmerian tectonic phase (see Van Adrichem Boogaert & Kouwe, 1997; De Jager, 2007)

Outside the Netherlands in Britain Late Permian or Early Triassic rifting have been proposed. In the Netherlands significant tectonic movements during the Triassic are concluded from three unconformities.

They are called the Hardegsen unconformity (also called Solling unconformity) found over large parts of the Netherlands separating the Upper and the Lower Germanic Trias groups. In the upper Germanic Trias group one can find the early Kimmerian unconformities in the Keuper. These unconformities are related to rifting. In the North Sea four rift grabens can be observed located in the Netherlands, Germany and Denmark. They are called the Dutch Central Graben, Horn Graben and the Glückstad Graben. The thickness of Upper Triassic and Lower Jurassic section can exceed 1500m. There is an increased thickness of that section in the Broad Fourteens basin. This may be related to the overall stress condition of the area but is not interpreted as rifting.

3.2.2.2 Mid and late Kimmerian tectonic phase

The major tectonic elements of the Netherlands form during the Mid and late Kimmerian tectonic phase in the second half of the Jurassic.

Early in the Middle Jurassic (Aalenian-Bathonian) a large area in the middle of the North Sea, the Central North Sea Dome, was uplifted by the mid Kimmerian tectonic movements (Ziegler, 1990, Underhill & Partington, 1993). Regional uplift and erosion over most of the Netherlands occurred during the Late Jurassic lasting in some areas till the Early Cretaceous. The doming is generally associated with a new phase in the Mesozoic break-up of the Pangea supercontinent between Greenland and Scandinavia. After the doming the well known trilete rift system developed in the northern North Sea, well north of the Dutch offshore. The southern branch called the Central Graben, is via Tail End and Feda half-grabens connected to the Dutch part of the North Sea via the German and Danish offshore (see: Wride, 1995).

The rough outlines of this Central North Sea rift dome are described by the erosional edge of the Early Jurassic series (Underhill & Partington, 1993). Similar large radius arches are associated with currently active volcanic rifts as the more recent Rhine Graben rift system.

The Central Graben transects the crest of this dome. At the triple junction of the Central Graben, the Viking Graben and the Moray Firth fault a large volcanic center was initiated. The presence of Permian salt complicates the structural interpretation. The Dutch Central Graben was probably the southernmost extension of a major N-S oriented depocenters, which extended from the Viking Graben into the Central North Sea Graben. Contrary to some of the other Jurassic graben structures is the Dutch Central Graben a reactivated Triassic rift structure. At the end of the Jurassic the rupturing of the Jurassic Dutch Central Graben reached the southern part of the Graben and was active in the Vlieland Basin (Herngreen et al., 1991). The marine incursions started in the Middle to Late Callovian (Clark-Lowes et al., 1987; Wong et al., 2007; -, 1989; Adrichem Boogaert & Kouwe, 1997). The Vlieland basin was flooded during the Portlandian. Rupturing started in the Broad Fourteens, West Netherlands, Central Netherlands and Roer Valley Graben during the Lower Cretaceous (see for general outline and stratigraphy: Adrichem Boogaert & Kouwe, 1997).

Towards the end of the Early Cretaceous crustal separation was in the north Atlantic and rifting began to concentrate on the area between Norway and Greenland (Ziegler, 1990). During the Early Cretaceous the continued rifting in the North Atlantic resulted in the formation of oceanic crust and Africa started to rotate anticlockwise towards the European plate. The South Atlantic started to open.

Consequently rifting in the Netherlands ceased and inversion started in the Jurassic basins perhaps as early as Albian (Oudmayer & De Jager, 1993)

It is interesting to note that the Mid- and Late-Kimmerian rifting did not cut through the old Caledonian and Lower Carboniferous highs and rejuvenated old basin areas like the Campine Basin.

3.2.3 The Alpine collision

The Alpine collision adds a compressional element to the rifting tectonics. Sissingh (2006) describes the repeatedly occurring changes in the plate collision stress regime between Europe, Africa, Iberia and Apulia, as well the stepwise opening of the North Atlantic and accompanying plate reorganization of the Mid-Atlantic Ridge. Old structural massifs like the Ardennes and the Rhenisch blocks are little affected. The basin areas show periodical deformation uplift and erosion (Sissingh, 2006),

Three major mainly compressive tectonic phases have been recognized and named in the North Sea rift system (De Jager, 2003; Ziegler, 1987).

These phases have been named:

- Sub Hercynian (Late Cretaceous)
- Laramide (Early Paleocene)
- Pyrenean (Eocene-Oligocene)

3.2.3.1 Subhercynian and Laramide late Cretaceous tectonic phases

In the late Cretaceous, subsidence patterns are completely reorganized. The previous stable highs started to subside rapidly, accumulating thick sequences of Chalk. Eventually even the long-lived London-Brabant Massif and the Mid North Sea and Ringkøbing-Fyn Highs were inundated.

Tectonic inversion affected many of the Late Jurassic-Early Cretaceous rift basins during Late Cretaceous and Early Tertiary. The sediments in the rift basins are more sensitive to compression and an inverted relief develops whereby graben areas are transformed in relative highs. See for inversion in the Netherlands (De Jager, 2007; De Jager, 2003; Gras & Geluk,

1999; Nalpas et al., 1995; Hooper et al., 1995; Gras, 1995; Baldschuhn et al., 1991; Dronkers & Mrozek, 1991). This change is governed by the progressive opening of the North Atlantic and the onset of the Alpine collision in the south in the realm of the western Tethys. The opening of the southern Atlantic causing an anticlockwise rotation and northwards movement of Africa changed the European stress pattern from extension to compression at the beginning of the late Cretaceous (Sissingh, 2006).

Locally, inversion as a result of sub-Hercynian compression tectonics, lead to significant erosion of the rift sequence. This erosion introduces considerable uncertainties in tracing the Late Cretaceous coastline corresponding to the maximum transgression.

Major erosion occurred during the various tectonic pulses. The hiatus in the Late Santonian to Early Campanian is called the sub-Hercynian unconformity. And can be found over several basins in the Netherlands and basins in the surrounding areas in Northwest Europe, like the Sole Pit, the West Netherlands Basin, the Lower Saxony Basin and the Dutch Central Graben Basin. The West Netherlands and the Broad Fourteens Basin display more inversion than the Dutch Central Graben.

The Laramide phase has been placed Mid-Paleocene (De Lugt, 2007) and terminated during the Late Paleocene. Further movements during the African-European collision result in compression of the whole European platform (Sissingh, 2006a) and are the formative stress source. The west-directed motion of Apulia characterizes this phase. Laramide compression caused inversion of the Dutch Central Graben, the Broad Fourteens Basin, the West Netherlands Basin and the UK Sole Pit Basin (De Jager, 2003; Ziegler, 1998; van Wijhe 1998). The Laramide phase was probably the most severe inversion (De Jager, 2003). Subsequent subsidence and rise of the area outside the North Sea basin caused a sedimentation pulse. Regional uplift caused massive erosion and peneplanation of complete areas.

Reactivation of faults after cessation of the Laramide phase resulted in the formation of local depocenters for example above the inverted Broad Fourteens Basin. It is observed that tectonic movement during the Paleocene was without exception accommodated by reactivation of Mesozoic and older faults. A dextral strike slip is suggested from structural geometries (Broad Fourteens Basin: Nalpas et al., 1995; Dutch Central Graben: Schroot, 1991).

3.2.3.2 Pyrenean and Savian tectonic phases

At the end of the Eocene the Pyrenean pulse caused broad uplift of the West and Central Netherlands basins. The Eocene-Early Oligocene Pyrenean orogenic phase resulted in renewed uplift (Ziegler, 1990; Van Wijhe, 1987). Sissingh (2006a) relates the movement to the slow north directed movement of Apulia that induces Alpine thrusting.

During this phase in the Netherlands inversion occurred and basin boundaries become overstepped and a large area compared to previous inversion pulses was severely uplifted. The aerial exposure of large parts of the basin results in serious erosion. The sediments are transported to the deeper parts of the basin. During the Pyrenean compressional phase strong variations occurred in accommodation space and erosion. The main stress orientation during the Pyrenean phase was at an oblique angle to the pre-existing structural grain, resulting in reactivation of faults with a strike slip component. This stress regime (Kooi & Cloetingh, 1989) is probably comparable to the earlier inversion phases but this is often obscured by the Pyrenean events. After the uplift almost comparable to the Laramide uplift, local subsidence is evident by Oligocene deposits that indicate differential subsidence.

A late tectonic pulse is called the Savian erosion. It is also believed to be caused by the regional stress regime induced by the Alpine orogeny (De Jager, 2007), but also sea-level changes may have played a role. The erosion terminates the Oligocene sedimentation. The event is clearly visible on seismic as the Mid-Miocene unconformity a mapable seismic marker widely recognized in the southern North Sea (Oudmayer & De Jager, 1993). Almost the whole southern North Sea emerged above sea level. In the late Oligocene Apulia moves slowly north resulting in crustal shortening in the Alpine areas this continues till the late Miocene when thrusting ends and also a re-organization of the North Atlantic sea floor spreading changes the stress regime.

After the Savian phase, regional subsidence resumed and delta sediments fill up the basin. No major tectonic movements can be seen in the sediment record. Present-day activity makes it likely that there was continuous movement in areas like the Roer valley Graben. The sediment thickness in these Grabens exceeds 2000m. Based also on observations of Dronkers & Mrozek (1991) and Nalpas et al. (1995) for Late Cretaceous-Early Tertiary movements, Dirkzwager et al. (2000) concludes that older faults in rifted basins have been reactivated numerous times. Basin boundaries have been reactivated in particular and seismicity along these boundaries illustrate that the rifting structures form a zone of structural weakness (Dirkzwager et al., 2000).

4.0 Fault patterns

For the understanding of the hydrocarbon trapping it is very important to understand the fault patterns. The most widely studied structural horizon in the Netherlands, the Base Zechstein, shows a typical rhomboid pattern of intersecting fault patterns (Oudmayer and De Jager, 1993; De Jager, 2007) with a dominant family of fault trends NW-SE, intersected by NNE-SSW faults. In the northern offshore a more N-S trend is found.

Interesting to note is that no significant difference is found in the rift basin area compared with the platform areas. Because the faults predate the rifting it leads to the conclusion that many faults are reactivated and few faults were newly formed with the exception of some pop up structures (De Jager, 2007; Glennie, 1998). Many faults at Rotliegend level were inherited from the fault pattern of the Late Carboniferous fault patterns that originated from the movements during the Variscan orogeny. Reactivation of faults can be very important for the understanding of reservoir properties like fault seal (*see Rijkers, this conference*).

Reactivation of faults makes it difficult to reconstruct paleo stress from present day fault patterns.

Shallower fault patterns are affected by salt movements because of the presence of Zechstein salts of large parts of the Netherlands. Salt walls have been found at the edges of the rift basins (Remmelts, 1996).

Inversion structures in the broad fourteens basin are more complex, with shorter wavelength, than in the Dutch Central graben, because the inversion was stronger (De Jager, 2003). The area of inversion is characterized by major reverse-faults.

5.0 Sedimentation and erosion in the Netherlands

Sedimentation in the Netherlands has been summarized by Ziegler (1982), Ziegler 1990 and Adrichem Boogaert & Kouwe (1997) and several chapters of Wong et al. (2007). For the sedimentation during the Permian and Triassic see also Geluk (2005). Detailed studies of the Tertiary (De Lugt, 2007) improve the understanding of the Tertiary events.

Because of significant erosion during the Jurassic uplift extrapolation outside the rift basins is often speculative.

Erosion during inversion periods is also difficult to estimate. A summary of the magnitude of inversion is given by De Jager (2003). In the Broad Fourteens crustal shortening has been estimated as 10-12%, whereas estimates for the Dutch Central Graben only reach 1-2%. The amount of uplift and erosion nowhere exceeds 2 km contrary to 3.5 km sometimes quoted.

5.1 Source rock and erosion

Understanding of the erosion is important to understand sediment loading important to understand hydrocarbon generation. A good understanding of uplift and erosion can be critical for the understanding of prospective areas (Alberts & Underhill, 1991). Erosion estimates are often very uncertain and sometimes controversial (De Jager & Geluk 2007) and is one of the least documented and understood subjects related to the rifting process.

For the Westphalian Limburg Formation coals, the most important gas generating source rock, the gas generation is closely interlinked with the Dutch rift basins. During the rifting phase gas generation accelerated in the subsiding rift basins and halted during inversion. Present day charge is limited to selected areas less affected by inversion (De Jager & Geluk 2007).

6.0 Magmatism related to rifting in the Netherlands

Rifting is closely linked to magmatism. The intensity of magmatism varies in the various rifting phases in the Netherlands.

- Extensive Permian Rotliegend volcanism in Germany, also to be found in the Netherlands (Van Bergen & Sissingh, 2007; Ziegler, 1992) may be linked to rifting.
- Evidence for Triassic magmatic activity is limited. Interesting are vulcanoclastics in the Central Graben well F3-7. Low level of Triassic volcanism has also been noted outside the Netherlands (Ziegler, 1982).
- More evidence for Jurassic magmatic activity can be found. The most significant feature is the Zuidwal Volcano (Cottençon et al., 1975; Herngreen et al. 1991). The Zuidwal volcanics may represent one or more cycles of trachite-phonolite eruptions (Van Bergen & Sissingh, 2007).

Other Jurassic magmatic rocks are present at the rim of the Dutch Central Graben (E6-1). These occurrences are considered the southernmost expression of Jurassic magmatism related to the rifting in the Northern North Sea with a centre in the Forties basaltic province.

- From the Cretaceous many magmatic occurrences have been described farther to the south in the West Netherlands Basin, illustrating that igneous activity becomes increasingly younger in southward direction.
- Ash layers in the Tertiary Dongen Formation and volcanic material in Oligocene strata illustrate that volcanic activity in the Tertiary is important. This is most likely linked to the development of the Rhine rift system (Van Bergen & Sissingh, 2007).

7.0 Heat flow in Graben areas

For the understanding of hydrocarbon generation heat flow is an important factor. In rift areas the heat flow is higher than in the surrounding areas. Based on 65 wells the average temperature gradient in the Netherlands is 31 degree C/km. Based on the subsurface values only, the trend is approximately 29.5 degree C/km (Verweij, 2003). The Roer valley Graben shows a higher gradient. In the Central Graben area from tests and logs a gradient of 31.7 – 35.5 degree C/km has been computed (Crepieux et al., 1996). Temperatures below the gradient are observed in northern blocks like A12, E17, F3, F18, L2, L8, L11, L12). In blocks P3, P5, P12, Q1, Q4 Q10 temperatures have been observed that exceed the trend.

8.0 Summary of the workshop

Structuration related to the rifting process affects large areas of the Dutch subsurface. Understanding of the processes related to rifting are important to understand structural style of hydrocarbon traps and also the generation of hydrocarbons in the Netherlands.

This summary shows that despite the vast increase of structural knowledge because of increased coverage of the on and offshore area by commercial seismic, the number of papers that discuss the Dutch rifted basin is limited. This overview shows that besides structural interpretation more work can be done on estimates of erosion and this literature survey outlines a few areas of special interest like the interpretation of deep seismic lines in the Netherlands.

We hope that this workshop will initiate more integrated studies of the rifting processes in the Netherlands.

9.0 Acknowledgments

I like to thank Energie Beheer Nederland B.V. for their support to publish this overview. Helpful discussions with Jan de Jager are gratefully acknowledged.

10.0 References

- Abramovitz, T., Landes, M., Thybo, H., Brian Jacob, A.W. & Prodehl, C. (1999)** *Crustal velocity structure across the Tornquist and Iapetus Suture Zones - a comparison based on MONA LISA and VARNET data.* Tectonophysics, vol. 314, p. 69-82.
- Abramovitz, T. & Thybo, H. (2000)** *Seismic images of Caledonian, lithosphere-scale collision structures in the southeastern North Sea along MONA LISA profile 2.* Tectonophysics, vol. 317, p. 27-54.
- Alberts, L.J.H., Geel, C.R. & Klasen, J.J. (2003)** *Reservoir characterisation using process-response simulations; the Lower Cretaceous Rijn Field, West Netherlands basin.* Geologie en Mijnbouw / Netherlands Journal of Geosciences, vol. 82, no. 4, p. 313-324.
- Alberts, M.A. & Underhill, J.R. (1991)** *The effect of Tertiary structuration on Permian gas prospectivity, Cleaver Bank area, southern North Sea, UK.* In: Spencer, A.M. (ed.). Generation, accumulation and production of Europe's hydrocarbons, Proceed. First Conf. EAPG. May 30 - June 2, West Berlin, Special Publication of the European Association of Petroleum Geoscientists, no. 1., Oxford University Press, p. 161-173.
- Baldschuhn, R., Best, G. & Kockel, F. (1991)** *Inversion tectonics in the north-west German basin.* In: Spencer, A.M. (ed.). Generation, accumulation and production of Europe's hydrocarbons, Proceed. First Conf. EAPG. May 30 - June 2, West Berlin, Special Publication of the European Association of Petroleum Geoscientists, no. 1., Oxford University Press, p. 149-159.
- Barnard, P.C. & Cooper, B.S. (1981)** *Oils and source rocks of the North Sea area.* In: Illing, L.V. & Hobson, G.D. (eds.). Petroleum geology of the Continental Shelf of North-West Europe: Proceedings of the 2nd Conference. Institute of Petroleum, London, p. 169-175.
- Bodenhausen, J.W.A. & Ott, W.F. (1981)** *Habitat of the Rijswijk oil province, onshore, the Netherlands.* In: Illing, L.V. & Hobson, G.D. (eds.). Petroleum geology of the Continental Shelf of North-West Europe: Proceedings of the 2nd Conference. Institute of Petroleum, London, p. 301-309.
- Cottencçon, A., Parant, B. & Flacelière, G. (1975)** *Lower Cretaceous Gas-Fields in Holland.* In: Woodland, A.W. (ed.). Petroleum and the Continental shelf of North-West Europe, vol. 1, Appl. Sc. Publishers LTD, Barking Essex, p. 403-412.
- Clark-Lowes D.D., Kuzemko, N.C.J. & Scott, D.A. (1987)** *Structure and petroleum prospectivity of the Dutch Central Graben and neighbouring platform areas.* In: Brooks, J. & Glennie, K.W. (eds.). Petroleum Geology of North-West Europe, Proceedings of the 3rd conference on Petroleum Geology of North West Europe, vol. 1, Graham and Trotman, London, p. 337-356.
- Crepieux, N., Sacleux, M. & Mathis, B. (1998)** *Influence of the pressure on the petroleum system. Example from Triassic in the Netherlands Central Graben.* In: Mitchel, A. & Grauls, D. (eds.). Overpressures in Petroleum Exploration. Proceedings workshop April, Pau, France, Elf-EP editions, Bulletin des Centres de Recherches Exploration-Production elf aquitaine, Mémoire 22, p. 123-132.
- De Jager, J., Doyle, M.A., Grantham, P.J. & Mabillard, J.E. (1996)** *Hydrocarbon habitat of the West Netherlands Basin.* In: Rondeel, H.E., Batjes, D.A.J. & Nieuwenhuijs, W.H. (eds.). Geology of gas and oil under the Netherlands. Royal Geological and Mining Society of the Netherlands, Kluwer Academic Publishers,

Rifting systems and its significance for hydrocarbon exploration in the Netherlands

Dordrecht, p. 191-209.

- De Jager, J. (2003)** *Inverted basins in the Netherlands, similarities and differences*. Geologie en Mijnbouw/ Netherlands Journal of Geosciences, vol. 82, no. 4, p. 339-349.
- De Jager, J. (2007)** *Geological development*. In: Wong, T.E., Batjes, D.A.J. & Jager, J. de (eds.). Geology of the Netherlands, Royal Netherlands Academy of Arts and Sciences, Amsterdam, p. 5-26.
- De Jager, J. & Geluk, M.C. (2007)** *Petroleum Geology*. In: Wong, T.E., Batjes, D.A.J. & Jager, J. de (eds.). Geology of the Netherlands, Royal Netherlands Academy of Arts and Sciences, Amsterdam, p. 241-264.
- De Jong, M.G.G. & Laker, N. (1992)** *Reservoir modeling of the Vlieland Sandstone of the Kotter Field (Block K18b), offshore, The Netherlands*. Geologie en Mijnbouw, vol. 71, p. 173-188.
- De Lugt, I.R. de (2007)** *Stratigraphical and structural setting of the Palaeogene siliciclastic sediments in the Dutch part of the North Sea Basin*. Geologica Ultraiectina, No. 270, Ph D. thesis University of Utrecht, 112 p.
- Dirkzwager, J.B., Van Wees, J.D.A.M., Cloetingh, S.A.P.L., Geluk, M.C., Dost, B. & Beekman, F. (2000)** *Geo-mechanical and rheological modelling of upper crustal faults and their near-surface expression in the Netherlands*. Global and Planetary Change, vol. 27, no. 1, p. 67-88.
- Dronkers, A.J. & Mrozek, F.J. (1991)** *Inverted basins of The Netherlands*. First Break, vol. 9, p. 409-425.
- Duin, E.J.T., Rijkers, R.H.B. & Remmelts, G. (1994)** *Deep seismic reflections in the Netherlands, an overview*. Geologie en Mijnbouw, vol. 64, p. 191-197.
- Duin, E.J.T., Doornenbal, J.C., Rijkers, R.H.B., Verbeek, J.W. & Wong, T.E. (2006)** *Subsurface structure of the Netherlands - results of recent onshore and offshore mapping*. Geologie en Mijnbouw / Netherlands Journal of Geosciences, vol. 85, no. 4, p. 245-276.
- Geluk, M.C. (1990)** *The Cenozoic Roer Valley Graben, southern Netherlands*. Mededelingen Rijks Geologische Dienst, vol. 44, p. 65-72.
- Geluk, M.C. (2005)** *Stratigraphy and tectonics of Permo-Triassic basins in the Netherlands and surrounding areas*. Ph.D. Thesis University of Utrecht, 171 p.
- Glennie, K.W. [ed.] (1998)** *Petroleum Geology of the North Sea, Basic concepts and recent advances*. Blackwell Science Ltd, London, 636 p.
- Goh, L.S. (1996)** *The Logger oil Field: Reservoir Architecture and Heterogeneity*. In: Rondeel, H.E., Batjes, D.A.J. & Nieuwenhuijs, W.H. (eds.). Geology of gas and oil under the Netherlands. Royal Geological and Mining Society of the Netherlands, Kluwer Academic Publishers, Dordrecht, p. 255-264.
- Gras, R. (1995)** *Late Cretaceous sedimentation and tectonic inversion, southern Netherlands*. Geologie en Mijnbouw, vol. 74, p. 117-127.
- Gras, R. & Geluk, M.C. (1999)** *Late Cretaceous-Early Tertiary sedimentation and tectonic inversion in the southern Netherlands*. Geologie en Mijnbouw, vol. 78, p. 1-19.
- Hastings, A., Murphy, P. & Stewart, L. (1991)** *A multi-disciplinary approach to reservoir characterization: Helm field, Dutch North Sea*. In: Spencer, A.M. (ed.). Generation, accumulation and production of Europe's hydrocarbons, Proceed. First Conf. EAPG. May 30 - June 2, West Berlin, Special Publication of the European Association of Petroleum Geoscientists, no. 1., Oxford University Press, p. 193-202.
- Herngreen, G.F.W., Smit, R. & Wong, T.E. (1991)** *Stratigraphy and tectonics of the Vlieland basin, The Netherlands*. In: Spencer, A.M. (ed.). Generation, accumulation and production of Europe's hydrocarbons, Proceed. First Conf. EAPG. May 30 - June 2, West Berlin, Special Publication of the European Association of Petroleum Geoscientists, no. 1., Oxford University Press, p. 175-192.
- Heybroek, P. [comp.] (1974)** *Explanation to tectonic maps of the Netherlands*. Geologie en Mijnbouw, vol. 53, no. 2, p. 43-50.
- Heybroek, P. (1975)** *On the structure of the Dutch part of the Central North Sea Graben*. In: Woodland, A.W. (ed.). Petroleum and the Continental shelf of North-West Europe, vol. 1, Appl. Sc. Publishers LTD, Barking Essex, p. 339-351.
- Hooper, R.J., Goh, L.S. & Dewey, F.J. (1995)** *The inversion history of the northeastern margin of the Broad Fourteens Basin*. In: Buchanan, J.G. & Buchanan, P.G. (eds.). Basin Inversion, The Geological Society of London, Special Publication, no. 88, p. 307-317.
- Kearey, P. & Vine, F.J. (1990)** *Global tectonics*. Blackwell Scientific Publications, Oxford, 302 p.

Rifting systems and its significance for hydrocarbon exploration in the Netherlands

- Kockel, F. (2002)** *Rifting processes in NW-Germany and the German North Sea Sector*. Geologie en Mijnbouw / Netherlands Journal of Geosciences, vol. 81, no. 2, p. 149-158.
- Kockel, F. (2003)** *Inversion structures in Central Europe - Expressions and reasons, an open discussion*. Geologie en Mijnbouw / Netherlands Journal of Geosciences, vol. 82, no. 4, p. 367-382.
- Kooi, H. & Cloetingh, S.A.P.L. (1989)** *Intraplate stresses and the tectono-stratigraphic evolution of the central North Sea*. In: Tankard, A.J. & H.R. Balkwill, H.R. (eds.). Extensional Tectonics and Stratigraphy of the North Atlantic Margins, American Association of Petroleum Geologists, Memoir 46, published jointly with Canadian Geological Foundation, p. 541-558.
- Nalpas, T., Douaran, S. le, Brun, J.P., Unternehr, P. & Richert, J.P. (1995)** *Inversion of the Broad Fourteens Basin, a small scale model investigation*. Sedimentary Geology, vol. 95, p. 237-250.
- Nalpas, T., Richert, J.P., Brun, J.P., Mulder, T. & Unternehr, P. (1996)** *Inversion du "Broad Fourteens Basin" ou Graben de La Haye (sud de la Mer du Nord); apports de la sismique 3D*. Bulletin des Centres de Recherches Exploration-Production elf aquitaine, vol. 20, p. 309-321.
- Oudmayer, B.C. & De Jager, J. (1993)** *Fault reactivation and oblique-slip in the Southern North Sea*. In: Parker, J.R. (ed.). Petroleum Geology of Northwest Europe: Proceedings of the 4th conference, Geological Society of London, vol. 2, p. 1281-1290.
- Remmelts, G. (1996)** *Salt tectonics in the southern North Sea, the Netherlands*. In: Rondeel, H.E., Batjes, D.A.J. & Nieuwenhuijs, W.H. (eds.). Geology of gas and oil under the Netherlands. Royal Geological and Mining Society of the Netherlands, Kluwer Academic Publishers, Dordrecht, p. 143-158.
- Rijkers, R.H.B., Duin, E.J.T., Duser, M. & Langenaeker, V. (1993)** *Crustal structure of the London Brabant Massif, southern North Sea*. Geological Magazine, vol. 130, p. 569-574.
- Rijkers, R.H.B. & Duin, E.J.T. (1994)** *Crustal observations beneath the southern North Sea and their tectonic and geologic implications*. Tectonophysics, vol. 240, p. 215-224.
- Roelofsen, J.W. & De Boer, W.D. (1991)** *Geology of the Lower Cretaceous Q/1 oil-fields, Broad Fourteens basin, The Netherlands*. In: Spencer, A.M. (ed.). Generation, accumulation and production of Europe's hydrocarbons, Proceed. First Conf. EAPG. May 30 - June 2, West Berlin, Special Publication of the European Association of Petroleum Geoscientists, no. 1., Oxford University Press, p. 203-216.
- Schroot, B.M. (1991)** *Structural development of the Dutch Central Graben*. In: Michelsen, O. & Frandsen, N. (eds.). The Jurassic in the Southern Central Trough, Danmarks Geologiske Undersøgelse series B, no. 16, p. 32-35.
- Sissingh, W. (2006)** *Kinematic sequence stratigraphy of the European Cenozoic Rift System and Alpine Foreland Basin: correlation with Mediterranean and Atlantic plate-boundary events*. Geologie en Mijnbouw / Netherlands Journal of Geosciences, vol. 85, no. 2, p. 77-129.
- Sissingh, W. (2006)** *Syn-kinematic palaeogeographic evolution of the West European Platform: correlation with Alpine plate collision and foreland deformation*. Geologie en Mijnbouw / Netherlands Journal of Geosciences, vol. 85, no. 2, p. 131-180.
- Underhill, J.R. & Partington, M.A. (1993)** *Jurassic thermal doming and deflation in the North Sea: implications of the sequence stratigraphic evidence*. In: Parker, J.R. (ed.). Petroleum Geology of Northwest Europe: Proceedings of the 4th conference, Geological Society of London, p. 337-245.
- Van Adrichem Boogaert, H.A. & Kouwe, W.F.P. (1997)** *Stratigraphic nomenclature of the Netherlands, revision and update by RGD and NOGPA*. Mededelingen Rijks Geologische Dienst, nieuwe serie, vol. 50, section a - j.
- Van Bergen, M.J. & Sissingh, W. (2007)** *Magmatism in the Netherlands: expression of the north-west European rifting history*. In: Wong, T.E., Batjes, D.A.J. & Jager, J. de (eds.). Geology of the Netherlands, Royal Netherlands Academy of Arts and Sciences, Amsterdam, p. 197-221.
- Van Wees, J.D.A.M., Stephenson, R.A., Ziegler, P.A., Bayer, U., McCann, T., Dadlez, R. Gaupp, R., Narkiewicz, M. Bitzer, F. & Scheck, M. (2000)** *On the origin of the Southern Permian Basin, Central Europe*. Marine and Petroleum Geology, vol. 17, nr. 1, p. 43-59. Wong, T.E., Batjes, D.A.J. & Jager, J. de [eds.] (2007) Geology of the Netherlands. Royal Netherlands Academy of Arts and Sciences, Amsterdam, 354 p.
- Verweij, J.M. (2003)** *Fluid flow systems analysis on geological timescales in onshore and offshore Netherlands*. With special reference to the Broad Fourteens Basin. Ph.D. Thesis VU, Amsterdam, 278 p.
- Verweij, J.M., Simmelink, H.J., Van Balen, R.T. & David, P. (2003)** *History of petroleum systems in the*

Rifting systems and its significance for hydrocarbon exploration in the Netherlands

southern part of the Broad Fourteens Basin. Geologie en Mijnbouw / Netherlands Journal of Geosciences, vol. 82, no. 1, Special issue 'on Geofluids in the Netherlands', p. 71-90.

- Visser, W.A., Van Dalen, W.J., Engelen, F.H.G. & Hillaert, J.G.A. (1987)** *The advance of geophysics in exploration*. In: Visser, W.A., Zonneveld, J.I.S. & Loon, A.J. van (eds.). Seventy-five years of geology and mining in The Netherlands (1912-1987). Royal Geol. and Mining Soc. of The Netherlands (KNGMG), The Hague, p. 167-198.
- Wong, T.E., Van Doorn, T.H.M. & Schroot, B.M. (1989)** *'Late Jurassic' petroleum geology of the Dutch Central North Sea Graben*. In: Poelchau, H.S. & Mann, U. (eds.). Geologic Modelling. Aspects of integrated basin analysis and numerical simulation. Geologische Rundschau, band 78, Heft 1, p. 319-336.
- Wong, T.E., Batjes, D.A.J. & Jager, J. de [eds.] (2007)** *Geology of the Netherlands*. Royal Netherlands Academy of Arts and Sciences, Amsterdam, 354 p.
- Wride, V.C. (1995)** *Structural features and structural styles from the Five Countries Area of the North Sea Central Graben*. First Break, vol. 13, no. 10, p. 395-407.
- Ziegler, P.A. (1975)** *North Sea Basin History in the Tectonic Framework of North-Western Europe*. In: Woodland, A.W. (ed.). Petroleum and the Continental shelf of North-West Europe, vol. 1, Appl. Sc. Publishers LTD, Barking Essex, p. 131-148.
- Ziegler, P.A. (1975)** *Geologic evolution of the North Sea and its tectonic framework*. American Association of Petroleum Geologists Bulletin, vol. 59, no. 7, p. 1073-1097.
- Ziegler, P.A. (1978)** *North-Western Europe: Tectonics and basin development*. Geologie en Mijnbouw, vol. 57, no. 4, p. 589-626.
- Ziegler, P.A. & Louwerens, C.J. (1979)** *Tectonics of the North Sea*. In: Oele, E., Schüttenhelm, R.T.E. & Wiggers, A.J. (eds.). The Quaternary history of the North Sea, Acta Universitatis Upsaliensis vol. 2, p. 7-22.
- Ziegler, P.A. (1981)** *Evolution of sedimentary basins in North-West Europe*. In: Illing, L.V. & Hobson, G.D. (eds.). Petroleum geology of the Continental Shelf of North-West Europe: Proceedings of the 2nd Conference. Institute of Petroleum, London, p. 3-39.
- Ziegler, P.A. (1982)** *Geological Atlas of Western and Central Europe*. Shell Internationale Petroleum Maatschappij B.V., Distributed by the Elsevier Scientific Publishing Company, 130 p.
- Ziegler, P.A. (1990)** *Geological Atlas of Western and Central Europe (2nd. Edition)*. Shell Internationale Petroleum Maatschappij B.V., Distributed by the Geological Society Publishing House, Bath, 239 p., 56 encl.
- Ziegler, P.A. (1992)** *North Sea rift system*. Tectonophysics, vol. 208, p. 55-75.
- Ziegler, P.A. (1994)** *Cenozoic rift system of western and central Europe: an overview*. Geologie en Mijnbouw, vol. 73, no. 2-4, p. 99-127.
- Ziegler, P.A. (2001)** *Dynamic processes controlling development of rifted basins*. EUCOR-URGENT Publication, no. 1, 51 p.